


THERMOPHILIC AND MESOPHILIC AMINOPEPTIDASES
FROM BACILLUS STEAROTHERMOPHILUS

H. Zuber and G. Roncari

Translation of "Thermophile und mesophile
Aminopeptidasen aus Bacillus stearothermophilus",
Angewandte Chemie, Vol. 79, No. 20, 1967,
pp. 906 - 907.

(NASA-TT-F-15901) THERMOPHILIC AND
MESOPHILIC AMINOPEPTIDASES FROM BACILLUS
STEAROTHERMOPHILUS (Scientific Translation
Service) 8 p HC \$4.00
CSCL 06M



| | | | |
|---|--|--|-----------|
| 1. Report No. NASA TT F-15,901 | 2. Government Accession No. | 3. Recipient's Catalog No. | |
| 4. Title and Subtitle Thermophilic and mesophilic amino-peptidases from Bacillus stearotherophilus | | 5. Report Date August 1974 | |
| | | 6. Performing Organization Code | |
| 7. Author(s) H. Zuber and G. Roncari | | 8. Performing Organization Report No. | |
| | | 10. Work Unit No. | |
| 9. Performing Organization Name and Address SCITRAN Box 5456 Santa Barbara, CA 93108 | | 11. Contract or Grant No. NASw-2483 | |
| | | 13. Type of Report and Period Covered Translation | |
| 12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546 | | 14. Sponsoring Agency Code | |
| 15. Supplementary Notes Translation of "Thermophile und mesophile Aminopeptidasen aus Bacillus stearotherophilus". Angewandte Chemie, Vol. 79, No. 20, 1967, pp. 906 - 907. | | | |
| 16. Abstract Various strains of B. stearotherophilus contain different proportions of three aminopeptidases. Obligately thermophilic strains contain more of the thermophilic enzyme; obligately mesophilic strains contain very little of it, and facultative strains contain similar amounts of the three. | | | |
| 17. Key Words (Selected by Author(s)) | | 18. Distribution Statement Unclassified - Unlimited | |
| 19. Security Classif. (of this report) Unclassified | 20. Security Classif. (of this page) Unclassified | 21. No. of Pages 6 | 22. Price |

THERMOPHILIC AND MESOPHILIC AMINOPEPTIDASES
FROM BACILLUS STEAROTHERMOPHILUS [1]

H. Zuber and G. Roncari*

We have been able to isolate three aminopeptidases, /906**
differing in molecular weight, thermal stability, and substrate
specificity, from several strains of *Bacillus stearothermophilus*
[2]. Aminopeptidase I (AP I) is thermostable and is, therefore,
designated as the thermophilic enzyme. Aminopeptidases II and
III (AP II and AP III) are mesophilic enzymes, and thermolabile.
All three enzymes are metalloenzymes. In the native form, they
may exist as Co^{2+} or Mn^{2+} enzymes, as Co^{2+} and Mn^{2+} reactivate
the enzymes best after inactivation with EDTA ($\text{Co}^{2+} = \text{Mn}^{2+} > \text{Mg}^{2+} >$
 $[\text{Ni}^{2+} > \text{Ca}^{2+}]$). Ca^{2+} , Zn^{2+} and Fe^{2+} do not activate, but inhibit
at concentrations from 0.001 to 0.05 M. The pH optimum is at
pH = 7.5 to 9 for all three enzymes. The substrate specificity
(Table 2) is in part sharply different from the specificity of
the leucine aminopeptidase (LAP) [3] from hog kidneys.

AP I has a specific activity of 1,000 units $\text{mg}^{-1} \cdot \text{min}^{-1}$,
if we define the unit as hydrolysis of 1 μmol of substrate per
minute. The molecular activity is 150,000 $\text{mol} \cdot (\text{mol Enzyme}^{-1}) \cdot \text{min}^{-1}$.
Both activities were determined by hydrolysis of a 0.045 M
solution of Gly-Leu-Tyr in 0.05 M Tris-HCl buffer, pH=7.2,
 10^{-3} M Co^{2+} at 65° C.

AP I was purified by repeated chromatography in 0.05 M
Tris-HCl buffer (10^{-3} M Co^{2+}) on Sephadex G-150 and DEAE-
Sephadex A-50. AP II and AP III were purified by repeated
chromatography on Sephadex G-100 and DEAE-Sephadex. AP I,

* Private lecturer Dr. H. Zuber and Dr. G. Roncari, Laboratory for
microbiology of the Swiss Federal Institute of Technology

** Numbers in margin indicate pagination in original foreign text.

Table 1. PROPERTIES OF THE AMINOPEPTIDASES I - III

| | AP I | AP II | AP III |
|---|--------------|---------------------|---------------------|
| Molecular weight ^a | [ca. 150000] | [ca. 70000] | [ca. 80000] |
| Michaelis constant, K_m (60°C, Leu-Gly)(mmol/l) | [95] | [37] | [4] |
| Activation energy ^b (kcal/mole) | [16.3] | [16.3] | |
| Optimum temperature ^b (°C) | [90] | [60] | [60] |
| Thermostability ^c (Percent of the initial activity after 3 min 18 min 120 min at 80°C) | | [64] [35] [0] | [81] [34] [0] |

^a Determined by gel filtration with Sephadex G-200, 0.05 M Tris-HCl buffer, pH = 7.2, 10^{-3} M Co^{2+} .

^b Determined through hydrolysis of a 0.0075 - 0.06 M solution of Leu-Gly (AP I and AP II) or a 0.0002 - 0.002 M solution of Leu-Gly (AP III) (0.05 M Tris-HCl buffer, pH = 7.2, 10^{-3} M Co^{2+} ; incubation time: 10 minutes at 20, 40, 60, 80, 90 °C).

^c Determined by hydrolysis of a 0.06 M Leu-Gly solution, 0.05 M Tris-HCl buffer, pH = 7.2, 10^{-3} M Co^{2+} ; incubation time: 10 minutes at 65°C (AP I) or 37°C (AP II and AP III).

AP II and AP III are homogeneous in polyacrylamide gel electrophoresis [4] (pH = 8.9, 45 minutes). The first experiments in the ultracentrifuge yielded for AP I:
 $S_{20} = 2.80 \cdot 10^{-13}$ and $D_{20} = 1.8 \cdot 10^{-7} \pm 0.16$ (symmetric bands).

Table 2. SUBSTRATE SPECIFICITIES

(Substrate concentration: 0.045 M, 65°C, 0.05 M Tris-HCl buffer, pH = 7.2, 0.001 M Co^{2+}). The numbers, except for column C, indicate percent hydrolysis in 5 minutes.

| Peptide | AP I | C_1^a | AP II | AP III |
|-------------|------|---------|-------|--------|
| Leu-Gly | 16 | 30 | 1,6 | 1,6 |
| Gly-Leu | 0,8 | 1,4 | 0,1 | 0,09 |
| Leu-amid | 5 | 8,8 | 9 | 1,2 |
| Leu-Gly-Gly | 30 | 50 | 25 | 80 |
| Gly-Leu-Tyr | 36 | 77 | 8 | 18 |
| Pro-Tyr-Lys | 3,2 | 5,6 | 2,7 | 25 |
| Glu-Ala-Ala | 5,5 | 9,8 | 1,2 | 1,7 |
| Lys-Tyr-Glu | 1,5 | 2,6 | 1,3 | 24 |

^a $C_1 = k_1/E$: k_1 = first order reaction rate; E = protein concentration (mg N/ml), assuming 15% nitrogen in the protein.

From this it appears that AP I exists in solution as a severely hydrolysed or very extended molecule. AP I is stable in 8 M urea solution, but is inactivated and split into fragments by reagents which break hydrogen bonds, such as formic acid or LiBr.

All three enzymes are found as early as the beginning of the log phase in the bacterial growth curve. They appear both in the cells and in the culture medium. The yield of the enzyme is maximal at the end of the log phase and in the stationary phase (24-30 hours). AP I, AP II and AP III are present in different amounts in the different strains of *Bacillus stearothermophilus*. Obligately thermophilic strains (temperature optimum at 55°C, no growth at 37°C) produce much AP I and AP II and little AP III, while obligately mesophilic strains

(temperature optimum 37°C, /no growth at 55°C) contain much AP III and very little AP I and AP II. Facultative strains (growth at 37°C and 55°C) can contain the three aminopeptidases in comparable amounts.

In order to obtain the crude enzyme mixture, the bacteria were disrupted with ultrasound or in the sand mill. Lysozyme treatment liberates only AP II and AP III. AP I appears to be bound more strongly to the cell membranes. The aminopeptidases were obtained from the culture medium by adsorption on DEAE-Sephadex (pH = 7.2) and elution with 1 M NaCl solution (0.05 M Tris-HCl buffer, pH = 7.2, 10^{-3} M Co^{2+}).

REFERENCES

1. This work was supported by the Swiss National Fund for Promotion of Scientific Research (Project No. 3426).
2. ATCC 7953, 7954; NC1B 8924, 8157, 8920, 8222
3. Smith, E. L. and D. H. Spackman. J. Biol. Chemistry Vol. 212, 1955, p. 271.
4. Davis, B. J. Ann. New York Acad. Sci. Vol. 121, 1964, p.404.

Translated for National Aeronautics and Space Administration under contract No. NASw 2483, by SCITRAN, P. O. Box 5456, Santa Barbara, California, 93108.